PASJ: Publ. Astron. Soc. Japan , 1–??,
© 2013. Astronomical Society of Japan.

arXiv:1304.6471v1 [astro-ph.SR] 24 Apr 2013

Decretion discs around rapidly rotating stars

Umin Lee
Astronomical Institute, Tohoku University, Sendai, Miyagi 980-8578, Japan
lee@astr.tohoku.ac.jp

(Received ; accepted)

Abstract

We discuss steady viscous Keplerian decretion discs around rapidly rotating stars. We assume that low frequency modes, which may be excited by the opacity bump mechanism, convective motion in the core, or tidal force if the star is in a binary system, can transport an enough amount of angular momentum to the region close to the stellar surface. Under this assumption, we construct a star-disc system, in which there forms a viscous decretion disc around a rapidly rotating star because of the angular momentum supply. We find a series of solutions of steady viscous decretion discs around a rapidly rotating star that extend to $R_{\rm disc} \gtrsim 10R_*$ with R_* being the equatorial radius of the star, depending on the amount of angular momentum supply. If the supply is low, the extension $R_{\rm disc}$ becomes small.

1. Introduction

Discs around Be stars are now believed to be viscous Keplerian decretion discs (e.g., Porter & Rivinius 2003; Lee, Saio, & Osaki 1991). Formation mechanism for such decretion discs have not yet been identified, but it has been suggested that angular momentum transferred by non-axisymmetric oscillation modes could play a role in the formation of viscous discs around rapidly rotating B stars (e.g., Ando 1983, 1986; Lee & Saio 1993). Angular momentum transfer by non-axisymmetric oscillation modes in rotating stars has been discussed by various authors. Circularization of binary orbit and synchronization between the spin of a massive star and the orbital motion of the companion in a binary system are believed to result from angular momentum exchange between the orbital motion and the spin of the stars, where angular momentum redistribution in the star takes place through dissipative low frequency g-modes tidally excited by the orbital motion of the companion (e.g., Zahn 1975, 1977; Goldreich & Nicholson 1989; Papaloizou & Savonije 1997; Witte & Savonije 2001; Willems, van Hoolst, & Smeyers 2003). The problem of angular momentum transport in the Sun by low frequency g-modes has long been a hot topic in the field since the early years of helioseismology and investigated by many researchers including Press (1981), Schatzman (1993), Gough (1997), Kumar, Talon, & Zhan (1999), Talon, Kumar, & Zahn (2002), Mathis et al. (2008), where low frequency modes are assumed to be excited by convection motion in the convective envelope and suffer radiative damping as they propagate into the radiative core.

To discuss angular momentum transfer by non-axisymmetric oscillations in a rotating star, we use a theory of wave-meanflow interaction, in which fluid motions are separated into waves and the mean flow and dissipative processes of the waves propagating in the mean flow have an essential role for the forcing on the meanflow (e.g., Andrews & McIntyre 1978ab; Dunkerton 1980; Grimshaw 1984; Ando 1983; Goldreich & Nicholson 1989). For rotating stars, we regard the rotation velocity field as the meanflow and global oscillations as waves. In the Cowling approximation, in which the Euler perturbation of the gravitational potential is neglected, the meanflow equation may be given by (see the Appendix)

$$\bar{\rho}\frac{d\left\langle\hat{\ell}\right\rangle}{dt} = \frac{m}{2}\mathrm{Im}\left[\nabla\cdot(\boldsymbol{\xi}^* p')\right], \tag{1}$$

where $\boldsymbol{\xi}$ and p' are respectively the displacement vector and the Eulerian pressure perturbation of an oscillation mode and $\hat{\ell} \equiv \hat{r}\sin\hat{\theta}\hat{v}_\phi$ is the specific angular momentum and $\left\langle\hat{\ell}\right\rangle = (2\pi)^{-1}\int_0^{2\pi}\hat{\ell}d\phi$ denotes its zonal average, and we have separated the coordinates $\hat{x}_\alpha$ into two parts such that $\hat{x}_\alpha = x_\alpha + \xi_\alpha(t,\boldsymbol{x})$, where x_α are the Lagrangian mean coordinates in the sense that $\langle\xi_\alpha\rangle = 0$ with ξ_α being the displacement vector. The right-hand-side of equation (1) represents the forcing effects associated with the oscillation mode, and an example of the forcing term calculated for low frequency modes in a slowly pulsating B star is given in the Appendix.

In this paper, we try to construct a disc-star model, which is composed of a rapidly rotating star and a decretion disc around it. To support a decretion disc around a rotating star an enough amount of angular momentum must be supplied to the surface layers of the star, and this angular momentum supply is assumed to result from angular momentum deposition by axisymmetric oscillation modes. The rotation velocity in the surface layers is accelerated acquiring angular momentum transferred by non-axisymmetric low frequency modes and a viscous Keplerian decretion disc forms around the star with the excessive angular momentum. Since the κ-mechanism associated with the iron opacity bump excites low frequency and high radial order g-modes and r-modes in slowly pulsating B (SPB) stars and low radial order f- and p-modes in β Cephei stars (Dziembowski et al. 1993, Gautschy & Saio 1993), we may use the low frequency modes as an agent which brings about angular momentum transfer for SPBe stars. For early Be stars, however, the opacity bump mechanism does not work for driving low frequency modes, and we may resort to stochastic excitation of low frequency modes by convective motion in the core (Neiner et al 2012). If the Be star is in a binary system, low frequency modes may be tidally excited by the orbital motion of the companion star and they could work for disc formation.

Okazaki (2001) has calculated steady and transonic decretion discs around Be stars, taking account of the effects of the radiation pressure on the radial flow. He has shown that the velocity field in the disc is very close to the Keplerian near the star and tends to angular momentum conserving in the region far from the star and that the effects of the radiation pressure are not essential for the existence of the transonic disc solutions. In his numerical analysis, he did not treat a rotating star and a disc around it as one system and assumed no angular momentum source in the subsurface layers of the star. These two points are what we concern in this paper, and we are interested in the part of the system around the rotating star.

In the angular momentum conservation equation for a disc-star system, we need to consider both the forcing term, which appears in the meanflow equation and may be regarded as a source of angular momentum in the subsurface layers of the star, and the term due to the viscous torque, which is essential for angular momentum transfer in the disc. We derive in §2 a set of ordinary differential equations for a disc-star system, in which a Keplerian viscous decretion disc is assumed to form around a rapidly rotating star. In §3, we give numerical results obtained for steady disc-star systems, and we conclude in §4. In the Appendix we derive the meanflow equation we use in this paper.

2. Equations for Decretion Discs around Rotating Stars

For rapidly rotating SPB stars, for example, numerous prograde sectoral g-modes and r-modes are destabilized by the opacity bump mechanism (e.g., Aprilia, Lee, & Saio 2012). Knowing their amplitudes and summing up all accelerating and decelerating contributions to the forcing on the meanflow, we can in principle estimate the amount of angular momentum deposited to the surface layers. With a linear theory of oscillations, however, we have no means to determine the amplitudes. In this paper, assuming that the contributions to accelerating the surface layers are dominant over those to deceleration, we consider a disc-star system, in which there forms around a rapidly rotating star a viscous Keplerian decretion disc, supported by excessive angular momentum transferred by the low frequency modes.

To treat both a rotating star and a viscous disc around it as one system, we work in cylindrical coordinates (R,φ,z). The displacement vector of an oscillation mode may be given by $\boldsymbol{\xi}=\xi_R\boldsymbol{e}_R+\xi_\varphi\boldsymbol{e}_\varphi+\xi_z\boldsymbol{e}_z$. If the amount of angular momentum (torque) provided by the low frequency modes is large enough to support a Keplerian viscous decretion disc around the star, the equation for angular momentum conservation, which includes both the viscous torque term for the disc and the forcing term due to the non-axisymmetric oscillations in the surface layers of the star, may be given by

$$\rho\frac{d}{dt}\left(Rv_\varphi\right)=\frac{1}{R}\frac{\partial}{\partial R}\left(R^2\sigma_{R\varphi}\right)+\frac{m}{2}\mathrm{Im}\left[\frac{1}{R}\frac{\partial}{\partial R}\left(R\xi_R^*p'\right)+\frac{\partial\left(\xi_z^*p'\right)}{\partial z}\right], \tag{2}$$

where $\sigma_{R\varphi}$ is the $R\varphi$ component of the viscous tensor and is assumed to dominate other components of the tensor for a geometrically thin disc. Here, we assume that the system is axisymmetric and in a steady state when the oscillation amplitudes are saturated, for example, by non-linear couplings between the oscillation modes (e.g., Lee 2012). If this is the case, integrating vertically equation (2), we obtain

$$\dot{M}\frac{\partial}{\partial R}\left(R^2\Omega\right)=\frac{\partial}{\partial R}\left(2\pi R^3\frac{\partial\Omega}{\partial R}\int_{-z_0}^{z_0}\eta dz\right)+\pi m\frac{\partial}{\partial R}R\int_{-z_0}^{z_0}dz\,\mathrm{Im}\left(\xi_R^*p'\right), \tag{3}$$

where we have used $\sigma_{R\varphi}=\eta R(\partial\Omega/\partial R)$ with η being the shear viscosity coefficient, $v_\varphi=R\Omega$, and we have assumed $p'=0$ at the surface given by $z=z_0(R)$. Here, $\dot{M}\equiv 2\pi R\Sigma v_R$ is the mass decretion rate, which is assumed a constant, and $\Sigma=\int_{-z_0}^{z_0}\rho dz$ is the surface density. Integrating equation (3) from R_0 to R, we obtain

$$2\pi R^3\frac{\partial\Omega}{\partial R}\int_{-z_0}^{z_0}\eta dz\equiv g(R)=\dot{M}j(R)-f(R)-\dot{J}, \tag{4}$$

where

$$j(R)=R^2\Omega, \tag{5}$$

$$f(R) = m\pi R \int_{-z_0}^{z_0} dz \,\mathrm{Im}\,(\xi_R^* p')\,, \tag{6}$$

and

$$\dot{J} \equiv \dot{M} j_0 - g_0 - f_0 = \dot{M} j - g - f, \tag{7}$$

and j_0, g_0, and f_0 denote the quantities evaluated at $R = R_0$.

Following Paczyńsky (1991), we also assume the disc-star system is in hydrostatic balance:

$$\frac{1}{\rho}\frac{\partial p}{\partial R} + \frac{\partial \Phi}{\partial R} = R\Omega^2, \tag{8}$$

$$\frac{1}{\rho}\frac{\partial p}{\partial z} + \frac{\partial \Phi}{\partial z} = 0, \tag{9}$$

where the gravitational potential is given by $\Phi = -GM_*/\sqrt{R^2 + z^2}$ with M_* being the mass of the star. Considering two neighboring points (R, z_0) and $(R + \delta R, z_0 + \delta z_0)$ on the surface of the disc-star system such that $p(R, z_0) = 0 = p(R + \delta R, z_0 + \delta z_0)$, we obtain, using equations (8) and (9) for hydrostatic balance,

$$\frac{dz_0}{dR} = \frac{R}{z_0}\left(\frac{r^3\Omega^2}{GM_*} - 1\right), \tag{10}$$

where $r = \sqrt{R^2 + z^2}$.

Rewriting equation (10) as $dr^2/dR^2 = r^3\Omega/GM_*$ and assuming Ω/GM_* is a constant, we integrate equation (10) to obtain

$$z_0^s(R) = \left[\left(\frac{1}{R_p} - \frac{\Omega^2 R^2}{2GM_*}\right)^{-2} - R^2\right]^{1/2}, \tag{11}$$

where $R_p = z_0^s(0)$ is the polar radius of the star. Using the condition $z_0^s(R_e) = 0$, we may define, as a function of R_p and Ω, the equatorial radius R_e of a star that is uniformly rotating at a rate Ω. We can also define the critical equatorial radius R_* of a star rotating at the critical rate $\Omega_0 \equiv (GM_*/R_*^3)^{1/2}$. For this critical radius, we have $R_*/R_p = 1.5$. Using this critical radius, we may rewrite (11) as

$$y_0(x) = \left[\left(\frac{1}{y_p} - \frac{\bar{\Omega}^2 x^2}{2}\right)^{-2} - x^2\right]^{1/2}, \tag{12}$$

where $x = R/R_*$, $y_0(x) = z_0^s(R)/R_*$, $y_p = R_p/R_* = 2/3$, and $\bar{\Omega} = \Omega/\Omega_0$. Note that $y_0(x) = 0$ gives the critical radius $x = 1$ at $\bar{\Omega} = 1$ for a uniformly rotating star. The rotation rate Ω may be normalized by $\Omega_e \equiv (GM_*/R_e^3)^{1/2}$, and in this case we have $\Omega/\Omega_e = x_e^{3/2}\bar{\Omega}$ with $x_e = R_e/R_*$, where x_e is determined as a solution to $y_0(x) = 0$ for a given $\bar{\Omega} \le 1$. Figure 1 plots x_e and Ω/Ω_e as a function of $\bar{\Omega} = \Omega/\Omega_0$. As $\bar{\Omega}$ decreases, x_e tends to $2/3$. We also note the rapid decrease of Ω/Ω_e as $\bar{\Omega}$ decreases from 1.

We divide the disc-star system into two parts, that is, the inner part ($R \le R_{\rm tr}$) and the outer part ($R \ge R_{\rm tr}$). We assume the inner part is uniformly rotating at a constant rate Ω_s and the surface of the inner part is given by equation (11) with $\Omega = \Omega_s$. On the other hand, the outer part of the system is composed of the outer part of the rotating star and a decretion disc and is allowed to rotate differentially. We assume the outer part extends to the radius $R_{\rm out} \gg R_*$.

For a thin disc, the shear viscosity coefficient η may be given as

$$\eta = \alpha \rho_0 v_{s,0} z_0, \tag{13}$$

where α is a dimensionless constant parameter such that $0 < \alpha < 1$, and ρ_0 and $v_{s,0} = (\partial P_0/\partial \rho_0)^{1/2}$ are the density and the sound speed evaluated at the equatorial plane. Assuming a polytropic relation $P = K\rho^{1+1/n}$, we integrate equation (9) to obtain

$$\rho = \rho_0 \left(1 - \frac{z^2}{z_0^2}\right)^n \quad \text{with} \quad \rho_0 = \left[\frac{GM_*}{2(n+1)KR^3}\right]^n z_0^{2n}, \tag{14}$$

and hence

$$v_{s,0} = \left(\frac{GM_*}{2n}\right)^{1/2} \frac{z_0}{R^{1.5}}, \tag{15}$$

and



Fig. 1. x_e (solid line) and Ω/Ω_e (dashed line) as a function of $\bar\Omega = \Omega/\Omega_0$, where $\Omega_e = (GM_*/R_e^3)^{3/2}$ and $\Omega_0 = (GM_*/R_*^3)^{3/2}$, and R_e and R_* are the equatorial radius of a star uniformly rotating at Ω and Ω_0, respectively, and M_* denotes the mass of the star.

$$\eta = \left[\alpha \frac{(GM_*)^{n+0.5}}{K^n}\right] \frac{z_0^{2n+2}}{R^{3n+1.5}}. \tag{16}$$

Equation (4) is now given by

$$g = \left[4\pi\alpha \frac{(GM_*)^{n+0.5}}{K^n}\right] \frac{z_0^{2n+3}}{R^{3n-1.5}} \frac{d\Omega}{dR} = \dot M R^2 \Omega - f - \dot J. \tag{17}$$

Equations (10) and (17) make a set of ordinary differential equations we have to solve with appropriate boundary conditions imposed at $R = R_{\rm tr}$ and $R = R_{\rm out}$.

In this paper, we assume $n = 1.5$ for the polytropic index. Equations (10) and (17) then reduce to

$$\frac{dy}{dx} = \frac{x}{y}\left[\bar\Omega^2 \left(x^2 + y^2\right)^{3/2} - 1\right], \tag{18}$$

$$\frac{d\bar\Omega}{dx} = \frac{1}{a}\frac{x^3}{y^6}\left[x^2\bar\Omega - b\hat f(x) - \dot j\right], \tag{19}$$

where

$$y = \frac{z_0(R)}{R_*}, \quad a = 4\pi\alpha\frac{G^2 M_*^2}{\dot M K^{1.5}}, \quad b = \frac{3}{4}\frac{M_*\Omega_0}{\dot M}, \quad \dot j = \frac{\dot J}{\dot M\sqrt{GM_* R_*}}, \tag{20}$$

and

$$\hat f(x) = x\int_{-y}^{y} \frac{dz}{R_*}\frac{\rho g r}{\bar\rho G M_*/R_*} x \sum_k m\,{\rm Im}\left(\frac{\xi_R^*}{r}\frac{p'}{\rho g r}\right), \tag{21}$$

and $\bar\rho = M_*/(4\pi R_*^3/3)$, and the summation in equation (21) is over the oscillation modes which contribute to the forcing on the meanflow, where k denotes a collective mode index. For the polytropic index $n = 1.5$, we may use $K = 0.4242 G M_*^{1/3} R_*$ (Chandrasekhar 1939) to obtain

$$a \simeq 4\pi\alpha\left(\frac{M_*}{\dot M}\right)\left(\frac{GM_*}{R_*^3}\right)^{1/2}, \tag{22}$$

and

$$\rho_0 \sim 1.5\bar\rho\ y^3/x^{9/2}. \tag{23}$$

Although the constant a depends on the viscosity parameter α, the magnitude of a is almost the same as that of the constant b, that is, $a \sim b$. As suggested by Paczyński (1991), the constant a can be as large as $a \sim 10^{12}\alpha$, depending on the quantities such as $\dot{M}$, M_*, and R_*. As a increases, however, it becomes difficult to numerically find solutions to the set of differential equations. In this paper, we employ $a = 10^7$, which leads to the disc thickness $z_0/R \sim 0.1$ (see below).

To determine the function $\hat{f}(x)$ with numerous low frequency non-adiabatic modes of a rapidly rotating star, we need to correctly estimate their amplitudes, which are difficult with a linear theory of oscillation. For our numerical computation, assuming that the resulting forcing function $\hat{f}(x)$ is that for accelerating the rotation velocity in the surface layers, we use a simple function given by

$$b\hat{f}(x) = b_0\left(\frac{1}{\exp\left[-b_1(x - x_0x_e)\right] + 1} - 1\right) \equiv b_0 f_0(x), \tag{24}$$

where b_0, b_1, and x_0 are parameters, and $x_e = R_e/R_*$. In the surface region of the star where $x \simeq x_e \leq 1$, we may approximate $|\hat{f}(x_1 \equiv x_0x_e)| \sim x_1^2 y_0(x_1)(\rho_0/\bar{\rho})|\sum_k m\mathrm{Im}[(\xi_R^*/r)(p'/\rho g r)]| \sim x_1^{-2.5} y_0^4(x_1)|\sum_k m f_k^2 \sin\delta_k|$ where $f_k \sim |\xi_R/r|$ is the normalized oscillation amplitude and δ_k is the phase difference between ξ_R and p' near the surface, and we have approximated $gr/(GM_*/R_*) \sim 1$. Since $|f_0(x_1)| = 0.5$, we may have $b_0 \sim b \times x_1^{-2.5} y_0^4(x_1)|\sum_k m f_k^2 \sin\delta_k|$. Although it is difficult to correctly estimate the magnitudes of the quantities such as $|\sum_k m f_k^2 \sin\delta_k|$, assuming $|\sum_k m \sin\delta_k| \sim 1$ and $x_1^{-2.5} y_0^4(x_1) \sim 10^{-3}$, we have $b_0 \sim 1$ for $f_k \sim 10^{-2}$ and $b \sim a = 10^7$. Note that for $x \gg 1$, we should have $b\hat{f}(x) = 0$, expecting there occurs no forcing on the meanflow in the disc.

The set of differential equations (18) and (19) are integrated with three boundary conditions, one given by $d\bar{\Omega}/dx = d\bar{\Omega}_K/dx$ with $\bar{\Omega}_K = x^{-3/2}$ at $x = x_{\rm out} \equiv R_{\rm out}/R_*$, and other two conditions by $\bar{\Omega}(x_{\rm tr}) = \bar{\Omega}_s$ and $y(x_{\rm tr}) = y_0(x_{\rm tr})$ at $x = x_{\rm tr} \equiv R_{\rm tr}/R_*$. Three boundary conditions for the second order ordinary differential equation are needed to determine the parameter $\dot{j}$. In this paper, we use $x_{\rm tr} = 0.8$.

As indicated by equation (19), for $x \gg 1$, $d\bar{\Omega}/dx$ changes its sign at x_j, for which $x_j^2\bar{\Omega}(x_j) - \dot{j} = 0$, where we have assumed $b\hat{f}(x) = 0$. If we assume $\bar{\Omega} \propto \bar{\Omega}_K$, we obtain $x_j \simeq \dot{j}^2$. To understand a rough property of the solution in the region $1 \ll x \ll x_j$ in which $x^2\bar{\Omega} \ll \dot{j}$, assuming functional forms given by $y = px^s$ and $\bar{\Omega} = q\bar{\Omega}_K$, where the parameters p, q, and s are assumed only weakly dependent on x, and substituting the expressions into equations (18) and (19), we obtain

$$s = \frac{11}{12}, \quad p \sim \left(\frac{\dot{j}}{aq}\right)^{1/6}, \quad q \sim \frac{1}{\left(1 + p^2x^{-1/6}\right)^{1/2}}. \tag{25}$$

These relations may be consistently satisfied if $p \sim 0.1$ and $q \sim 1$ for $\dot{j} \sim 10$ and $a \sim 10^7$, which indicates that the decretion discs are geometrically thin for the parameter values. If the disc extends beyond x_j, it may become angular momentum conserving so that $x^2\bar{\Omega} - \dot{j}$ be a constant (see Okazaki 2001). If this is the case, the outer boundary condition must be modified and an appropriate treatment of solutions around a point of x_j will be required.

3. Numerical Results

For the case of $\bar{\Omega}_s > 1$, we integrate equations (18) and (19) using a Henyey type relaxation method with the initial guesses given by $y = x[(\dot{j}x^{-1/2} - 1)/1.5a]^{1/6}$ and $\bar{\Omega} = x^{-3/2}$, and find the value of $\dot{j}$ that satisfies the three boundary conditions. For the case of $\bar{\Omega}_s \leq 1$, since there appears near $x \simeq 1$ a point x_m, at which $d\bar{\Omega}/dx = 0$, we integrate, for a given value of $\dot{j}$, the differential equations (18) and (19), with an implicit Runge-Kutta method, from $x = x_{\rm tr}$ using $y(x_{\rm tr}) = y_0(x_{\rm tr})$ and $\bar{\Omega}(x_{\rm tr}) = \bar{\Omega}_s$, to the point x_m. Since this integration gives x_m, $y_m = y(x_m)$, and $\bar{\Omega}_m = \bar{\Omega}(x_m)$ as a function of $\dot{j}$, or equivalently, y_m, $\bar{\Omega}_m$, and $\dot{j}$ as a function of x_m, we solve equations (18) and (19) in the interval between x_m and $x_{\rm out}$ with the relaxation method using the initial guesses to find the value of x_m such that the boundary conditions $y(x_m) = y_m$ and $\bar{\Omega}(x_m) = \bar{\Omega}_m$ at $x = x_m$ and $d\bar{\Omega}/dx = d\bar{\Omega}_K/dx$ at $x = x_{\rm out}$ are satisfied.

Figure 2 shows y and $\bar{\Omega}$ as a function of x for $\bar{\Omega}_s = 0.98$, 0.99, 1.00, and 1.01, where we have assumed $b_0 = b_1 = 50$, $x_{\rm out} = 10$, and $x_0 = 0.95$. In Table 1, we tabulate several characteristic quantities such as x_m and $\dot{j}$ as a function of $\bar{\Omega}_s$. As shown by the left panel of the figure, there appears a sharp dip in y at the boundary between the star and disc when $\bar{\Omega}_s < 1$, and the dip becomes deeper for smaller values of $\bar{\Omega}_s$. Table 1 indicates that this star-disc boundary is located at a radius near x_e for $\bar{\Omega}_s < 1$. For the case of $\bar{\Omega}_s \lesssim 1$, if we go outwards from $x_{\rm tr}$, at a point near x_e $\bar{\Omega}$ starts to rapidly increase to attain a super-Keplerian rate ($\Omega > \Omega_K$) at x_m and then decreases to the Keplerian velocity. Note that Ω is slightly sub-Keplerian in the region of $x \gtrsim 1$. However, for $\bar{\Omega}_s = 1.01$, we have $\Omega < \Omega_K$ in the entire x region of the solution and there appear no points of $d\bar{\Omega}/dx = 0$. For a given b_0, there exists the lower limit of Ω_s, below which no solutions to the differential equations are found. As $\bar{\Omega}_s$ decreases from 1, the amount of angular momentum deposition required to accelerate the sub-Keplerian rotation velocity to a super-Keplerian one is increased, and hence

Table 1. x_m and $\dot{j}$ for decretion disc solutions

Ω_s/Ω_0	x_e	Ω_s/Ω_e	x_m	$\dot{j}$
	$b_0=50$	$x_0=0.95$	$x_{\rm out}=10$	
0.9750	0.890	0.819	0.893	5.211
0.9800	0.900	0.837	0.905	4.859
0.9850	0.912	0.858	0.919	4.405
0.9900	0.926	0.883	0.936	3.837
0.9950	0.946	0.916	0.957	3.603
0.9980	0.965	0.946	0.963	5.420
0.9990	0.975	0.962	0.964	7.629
1.0000	1.000	1.000	0.965	16.81
1.0100	⋯	⋯	⋯	52.98
	$b_0=10$	$x_0=0.95$	$x_{\rm out}=2$	
0.9935	0.940	0.905	0.945	1.688
0.9940	0.942	0.908	0.948	1.666
0.9950	0.946	0.916	0.953	1.619
0.9960	0.952	0.925	0.960	1.578
0.9970	0.958	0.935	0.966	1.552
0.9980	0.965	0.946	0.972	1.577
0.9990	0.975	0.962	0.976	1.761
1.0000	1.000	1.000	0.976	3.160
1.0100	⋯	⋯	⋯	37.33
	$b_0=50$	$x_0=0.99$	$x_{\rm out}=10$	
0.9500	0.854	0.750	0.856	19.41
0.9600	0.867	0.774	0.870	18.77
0.9700	0.881	0.803	0.886	17.82
0.9800	0.900	0.837	0.907	16.39
0.9900	0.926	0.883	0.936	15.10
0.9950	0.946	0.916	0.952	16.86
0.9980	0.965	0.946	0.962	21.97
0.9990	0.975	0.962	0.966	25.70
1.0000	1.000	1.000	0.973	36.16
1.0100	⋯	⋯	⋯	66.14

the derivative $d\bar{\Omega}/dx$ inevitably becomes steeper in the region where acceleration takes place. The lower limit of $\bar{\Omega}_s$ is determined by the value at which $d\bar{\Omega}/dx=\infty$ or $y=0$. For $b_0=50$ and $x_{\rm out}=10$, the lower limit of $\bar{\Omega}_s$ is $\simeq 0.975$.

Figure 3 shows that the ratio y/x is less than ~ 0.1 in the disc, indicating the disc is geometrically thin. Since the ratio is approximately proportional to $\dot{j}^{1/6}$ as suggested by equation (25) and the value of $\dot{j}$ for $\bar{\Omega}_s=0.98$ is larger than that for $\bar{\Omega}_s=0.99$ (see Table 1), the ratio y/x for the former is larger than that for the latter. This figure also shows that the ratio gradually decreases as x increases from $x\sim 1$.

As suggested by Table 1, proper solutions to the differential equations can be obtained only when the outer boundary condition is imposed at $x_{\rm out} \lesssim x_j \simeq \dot{j}^2$, and the properties of the solutions for a given $\dot{j}$ do not strongly depend on $x_{\rm out}$ so long as $x_{\rm out} \lesssim x_j$ for $x_{\rm out}>x_m$. If we employ $b_0=10$ instead of $b_0=50$, we can find solutions for $x_{\rm out}=2$ but no solutions for $x_{\rm out}=10$ since the parameter value of $\dot{j}$ we obtain for $b_0=10$ is ~ 1.5 for $\bar{\Omega}_s<1$. The lower limit of $\bar{\Omega}_s$ for $b_0=10$ is $\simeq 0.9935$, which is much closer to unity than the lower limit 0.975 for the case of $b_0=50$. Figure 4 plots y and $\bar{\Omega}$ as a function of x for $b_0=10$. The discs for $b_0=10$ are thinner than for $b_0=50$, and the peak value of $\bar{\Omega}$ approximately corresponding to the lower limit of $\bar{\Omega}_s$ is smaller than that for the case of $b_0=50$.

As $\bar{\Omega}_s$ approaches unity from $\bar{\Omega}_s<1$, the parameter value $\dot{j}$ begins to increase rapidly (see Table 1). The large $\dot{j}$ means that decretion discs having large radii are possible, for example, a disc can have an extension of $x_{\rm out} \gtrsim 100$ if $\dot{j} \gtrsim 10$.

Assuming that the acceleration takes place in the region very close to the stellar surface, that is, in the region of much lower density, we examine the case of $x_0=0.99$, where we use $b_0=b_1=50$ and $x_{\rm out}=10$. We obtain a series of decretion disc solutions, the properties of which are quite similar to those for $x_0=0.95$, except for that the values of $\dot{j}$ for $x_0=0.99$ are greater than those for $x_0=0.95$. Because of the large values of the parameter $\dot{j}$, the discs can have a larger extension for $x_0=0.99$ than for $x_0=0.95$. The lower limit of $\bar{\Omega}_s$ is $\simeq 0.95$, which is smaller than $\simeq 0.975$ for the case of $x_0=0.95$.



Fig. 2. $y = z_0(R)/R_*$ (left panel) and $\bar{\Omega} = \Omega/\Omega_0$ (right panel) as a function of $x = R/R_*$ for $\bar{\Omega}_s = 0.98$, 0.99, 1.00, and 1.01 for $b_0 = b_1 = 50$, $x_0 = 0.95$, and $x_{\rm out} = 10$, where $\Omega_0 = (GM_*/R_*^3)^{3/2}$ with M_* being the mass of the star, and $\bar{\Omega}_s$ denotes the value of $\bar{\Omega}$ at $x = x_{\rm tr} = 0.8$. The dotted line in the right panel indicates the Keplerian rotation rate given by $\bar{\Omega}_K = x^{-3/2}$.



Fig. 3. The ratio y/x versus $x = R/R_*$ for $\bar{\Omega}_s = 0.98$, 0.99, 1.00, and 1.01 for $b_0 = b_1 = 50$ and $x_0 = 0.95$.



Fig. 4. $y = z_0(R)/R_*$ (left panel) and $\bar{\Omega} = \Omega/\Omega_0$ (right panel) as a function of $x = R/R_*$ for $\bar{\Omega}_s = 0.994$, 0.996, 0.998, 1.00, and 1.01 for $b_0 = 10$, $b_1 = 50$, $x_0 = 0.95$, and $x_{\rm out} = 2$. The dotted line in the right panel indicates the Keplerian rotation rate given by $\bar{\Omega}_K = x^{-3/2}$.



Fig. 5. $y = z_0(R)/R_*$ (left panel) and $\bar{\Omega} = \Omega/\Omega_0$ (right panel) as a function of $x = R/R_*$ for $\bar{\Omega}_s = 0.95$, 0.96, 0.97, 0.98, 0.99, 1.00, and 1.01 for $b_0 = b_1 = 50$, $x_{\rm out} = 10$, and $x_0 = 0.99$. The dotted line in the right panel indicates the Keplerian rotation rate given by $\bar{\Omega}_K = x^{-3/2}$.

4. Conclusion

We have obtained a series of solutions for steady viscous Keplerian decretion discs around a rapidly rotating star, assuming the existence of angular momentum supply to the region close to the surface of the star. The angular momentum supply may be provided by angular momentum deposition that takes place through wave-meanflow interaction, where the waves are low frequency global oscillations excited by the opacity bump mechanism for SPB star, or by a stochastic mechanism for early Be stars, or by a tidal force if the star is in a binary system. We may conclude that the angular momentum supply to the surface layers can be a mechanism for disc formation around rapidly rotating Be stars.

If the amount of angular momentum supply, which may be represented by the parameter b_0 in this paper, is large enough, viscous decretion discs can extend to a distance as far as $R_{\rm out} \gtrsim 10R_*$, but the possible disc radius may decrease as the supply becomes small. We also find that if the acceleration occurs in the region very close to the stellar surface, the possible extension a decretion disc attain can be as large as $R_{\rm out} \gtrsim 100R_*$. For a given b_0, there exists the lower limit of $\bar{\Omega}_s$, below which no steady decretion disc solutions are found. If b_0 is large enough, decretion disc solutions can be obtained even for the rotation rate $\sim$80% of the critical rate Ω_e determined by the equatorial radius R_e of the star, and the disc radius can be large enough for a Be star. If b_0 is small, however, disc solutions can be found only when $\bar{\Omega}_s$ is very close to unity. If the angular momentum supply is provided by global oscillations, the parameter b_0 is closely related to the amplitudes of the oscillations. We have argued that the amplitudes of order of $\xi_R/R_* \sim 0.01-0.1$ can lead to reasonable values of b_0.

The density at the mid-plane of the disc may be estimated by using equation (23), which leads to $\rho_0 \propto x^{-1.5}$ if the ratio y/x is assumed almost constant. Since $\rho_0 \propto x^{-3.5}$ has been suggested observationally, the x dependence of ρ_0 in our model is in a serious conflict with the observational estimation (e.g., Porter & Rivinius 2003). For B type stars we have $\bar{\rho} \sim 10^{-2}$, and since $y/x \sim 0.01-0.1$, we obtain $\rho_0 \sim 10^{-9}-10^{-6}$ at $x \sim 10$, the value of which is much higher than that observationally estimated (e.g., Waters 1986). We may be able to use much larger values for the parameter a to reduce the ratio y/x, but it becomes numerically difficult to obtain disc solutions to the differential equations.

As indicated by the plots of $\bar{\Omega}(R)$, there occurs a strong differential rotation in the region close to the surface, particularly for lower values of $\bar{\Omega}_s$. The strong differential rotation can change the modal properties of oscillations and hence the contributing to the forcing on the velocity field. Stability analysis of low frequency modes in differentially rotating star, which will be one of our future studies, is necessary if for the forcing we resort to the modes that are excited by the opacity bump mechanism.

Appendix 1. Mean Flow Equation in the Lagrangian Mean Formalism

Following Grimshaw (1984), in this Appendix we derive a meanflow equation for zonal flows around the rotation axis of stars, using the Lagrangian mean formalism (see also Andrews & McIntyre 1978b). In a frame rotating with the angular velocity Ω_c, the ϕ component of the momentum conservation equation in spherical polar coordinates $\hat{\boldsymbol{x}} \equiv (\hat{x}_1, \hat{x}_2, \hat{x}_3) = \left(\hat{r}, \hat{\theta}, \hat{\phi}\right)$ is given by

$$\hat{\rho}\left[\frac{d\hat{v}_\phi}{dt} + \frac{\hat{v}_r\hat{v}_\phi}{\hat{r}} + \frac{\hat{v}_\theta\hat{v}_\phi\cot\hat{\theta}}{\hat{r}} + 2\Omega_c\left(\sin\hat{\theta}\hat{v}_r + \cos\hat{\theta}\hat{v}_\theta\right)\right] = -\frac{1}{\hat{r}\sin\hat{\theta}}\frac{\partial\hat{p}}{\partial\hat{\phi}} - \hat{\rho}\frac{1}{\hat{r}\sin\hat{\theta}}\frac{\partial\hat{\Phi}}{\partial\hat{\phi}}, \tag{A1}$$

where

$$\hat{v}_r = \frac{d\hat{r}}{dt}, \quad \hat{v}_\theta = \hat{r}\frac{d\hat{\theta}}{dt}, \quad \hat{v}_\phi = \hat{r}\sin\hat{\theta}\frac{d\hat{\phi}}{dt}, \tag{A2}$$

and

$$\frac{d}{dt} = \frac{\partial}{\partial t} + \hat{v}_r\frac{\partial}{\partial\hat{r}} + \frac{\hat{v}_\theta}{\hat{r}}\frac{\partial}{\partial\hat{\theta}} + \frac{\hat{v}_\phi}{\hat{r}\sin\hat{\theta}}\frac{\partial}{\partial\hat{\phi}}. \tag{A3}$$

To discuss wave-mean flow interactions, we introduce the Lagrangian mean coordinates $\boldsymbol{x} \equiv (x_1, x_2, x_3) = (r, \theta, \phi)$ and the displacement vector ξ_α for $\alpha = 1,\ 2,\ 3$ such that

$$\hat{x}_\alpha = x_\alpha + \xi_\alpha(t, \boldsymbol{x}). \tag{A4}$$

We assume that for any given $\hat{v}_\alpha$ there is a unique "reference" velocity $\bar{v}_\alpha$, such that when the point x_α moves with velocity $\bar{v}_\alpha$ the point $\hat{x}_\alpha$ moves with velocity $\hat{v}_\alpha$ (e.g., Grimshaw 1984). Using the reference velocity, we may define

$$\frac{d}{dt} = \frac{\partial}{\partial t} + \bar{v}_r\frac{\partial}{\partial r} + \frac{\bar{v}_\theta}{r}\frac{\partial}{\partial\theta} + \frac{\bar{v}_\phi}{r\sin\theta}\frac{\partial}{\partial\phi}, \tag{A5}$$

where $\partial/\partial t$ indicates the partial time derivative with the coordinates $\boldsymbol{x}$ being held constant. For the wave-mean flow interaction formulatoin for zonal flows, we introduce an averaging procedure defined by

$$\langle f\rangle \equiv \frac{1}{2\pi}\int_0^{2\pi} f d\phi, \tag{A6}$$

and we assume for the displacement ξ_α

$$\langle \xi_\alpha\rangle = 0, \tag{A7}$$

where $\phi = x_3$. The velocity $\bar{\boldsymbol{v}}$ may be regarded as the mean velocity associated with the coordinates (x_α), and the displacement ξ_α represents the waves.

Using the Jacobian J for the coordinate transformation between $(\hat{x}_\alpha)$ and (x_α) given by

$$J \equiv \det\left(\frac{\partial \hat{x}_\alpha}{\partial x_\beta}\right) = \det\left(\delta_{\alpha\beta} + \frac{\partial \xi_\alpha}{\partial x_\beta}\right), \tag{A8}$$

we define the mean density $\tilde{\rho}$ associated with the coordinate $\boldsymbol{x}$ as

$$\hat{\rho}\hat{r}^2 \sin\hat{\theta} J = \tilde{\rho} r^2 \sin\theta. \tag{A9}$$

It is convenient to introduce (e.g., Andrews & McIntyre 1978b)

$$K_{\alpha\beta} = \frac{\partial J}{\partial\left(\partial \hat{x}_\alpha/\partial x_\beta\right)} = \frac{1}{2}\epsilon_{\alpha\kappa\lambda}\epsilon_{\beta\sigma\tau}\frac{\partial \hat{x}_\kappa}{\partial x_\sigma}\frac{\partial \hat{x}_\lambda}{\partial x_\tau}, \tag{A10}$$

for which we have

$$\frac{\partial K_{\alpha\beta}}{\partial x_\beta} = 0, \tag{A11}$$

and

$$K_{\alpha\beta}\frac{\partial \hat{x}_\alpha}{\partial x_\gamma} = \delta_{\beta\gamma} J = K_{\beta\alpha}\frac{\partial \hat{x}_\gamma}{\partial x_\alpha}, \tag{A12}$$

where we have employed the dual summation convention that Greek indices are summed over the range 1 to 3, and $\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1 = -\epsilon_{132} = -\epsilon_{213} = -\epsilon_{321}$ and $\epsilon_{\alpha\beta\gamma} = 0$ otherwise.

We rewrite equation (A1) as

$$\hat{\rho}\frac{d\hat{\ell}}{dt} = -\frac{\partial}{\partial\hat{\phi}}\hat{p} - \hat{\rho}\frac{\partial}{\partial\hat{\phi}}\hat{\Phi}, \tag{A13}$$

where $\hat{\ell}$ is the specific angular momentum, in an inertial frame, around the rotation axis defined by

$$\hat{\ell} = \hat{r}\sin\hat{\theta}\hat{v}_\phi + \Omega_c \hat{r}^2\sin^2\hat{\theta}. \tag{A14}$$

Multiplying equation (A13) by $\hat{r}^2\sin\hat{\theta}J$, we obtain

$$\tilde{\rho}\frac{d\hat{\ell}}{dt} + \frac{\partial\tilde{p}}{\partial\phi} = -\frac{1}{r^2\sin\theta}\frac{\partial}{\partial x_\beta}R_{3\beta} - \tilde{\rho}\frac{\partial}{\partial\hat{\phi}}\hat{\Phi}, \tag{A15}$$

where

$$R_{3\beta} = \delta_{3\beta}\left(J\hat{r}^2\sin\hat{\theta}\hat{p} - r^2\sin\theta\tilde{p}\right) - \hat{r}^2\sin\hat{\theta}\hat{p}\frac{\partial\xi_\gamma}{\partial x_3}K_{\gamma\beta}, \tag{A16}$$

and $\tilde{p} = \hat{p}(\tilde{\rho}, \hat{s})$ with $\hat{s}$ being the specific entropy, and we have used the identity

$$K_{3\beta}\hat{r}^2\sin\hat{\theta}\hat{p} = \delta_{3\beta}J\hat{r}^2\sin\hat{\theta}\hat{p} - \hat{r}^2\sin\hat{\theta}\hat{p}\frac{\partial\xi_\gamma}{\partial x_3}K_{\gamma\beta}. \tag{A17}$$

Applying the averaging procedure (A6) to equation (A15), we get

$$\tilde{\rho}\frac{d\left\langle\hat{\ell}\right\rangle}{dt} = -\frac{1}{r^2\sin\theta}\frac{\partial}{\partial x_\beta}\langle R_{3\beta}\rangle - \left\langle\tilde{\rho}\frac{\partial}{\partial\hat{\phi}}\hat{\Phi}\right\rangle, \tag{A18}$$

where we have used $\langle\partial\tilde{p}/\partial\phi\rangle = 0$. In general,

$$\langle R_{\alpha\beta}\rangle = \delta_{\alpha\beta}\left\langle J\hat{r}^2\sin\hat{\theta}\hat{p} - r^2\sin\theta\tilde{p}\right\rangle - \left\langle\hat{r}^2\sin\hat{\theta}\hat{p}\frac{\partial\xi_\gamma}{\partial x_\alpha}K_{\gamma\beta}\right\rangle \tag{A19}$$

is called the radiation stress tensor (e.g., Grimshaw 1984).

Equation (A18) may be regarded as the ϕ component of the meanflow equation, the left-hand-side of which may represent the time evolution of the meanflow and the right-hand-side the forcing by the waves represented by ξ_α. So far we have not assumed that the amplitudes of the waves ξ_α are infinitesimally small, and in principle we can formulate the wave-meanflow interaction as a nonlinear theory, which includes equations of motion for both the meanflows and waves. To avoid solving such a difficult non-linear problem, we use a linear theory to describe waves ξ_α and we calculate the forcing terms in the meanflow equation using the linear waves ξ_α.

If we employ a linear theory to describe waves represented by the displacement ξ_α, we may write

$$\hat{p} = \bar{p} + \delta p = \bar{p} + p' + \xi_\alpha \frac{\partial \bar{p}}{\partial x_\alpha}, \tag{A20}$$

where $\bar{p}$ is the pressure in equilibrium state, δp and p' denote the Lagrangian and Eulerian perturbations, respectively. Applying the averaging procedure (A6), to second order of the perturbations we obtain after some manipulations for the second term on the right hand side of equation (A16)

$$\left\langle \hat{r}^2 \sin\hat{\theta}\,\hat{p}\frac{\partial \xi_\gamma}{\partial x_3} K_{\gamma\alpha} \right\rangle = \left\langle r^2 \sin\theta \frac{\partial \xi_\alpha}{\partial x_3} p' \right\rangle + \left\langle \frac{\partial}{\partial x_\gamma}\left(r^2 \sin\theta \frac{\partial \xi_\alpha}{\partial x_3}\xi_\gamma \bar{p}\right)\right\rangle, \tag{A21}$$

where we have used $\langle \partial f/\partial\phi\rangle = \partial\langle f\rangle/\partial\phi = 0$ and $\partial\bar{p}/\partial\phi = 0$ for the equilibrium pressure $\bar{p}$, and $x_3 = \phi$. Because

$$\begin{aligned}\frac{\partial}{\partial x_\alpha}\frac{\partial}{\partial x_\gamma}\left(\bar{p}r^2\sin\theta\frac{\partial\xi_\alpha}{\partial\phi}\xi_\gamma\right) = {} & \frac{\partial}{\partial\phi}\left[\frac{\partial}{\partial x_\alpha}\frac{\partial}{\partial x_\gamma}\left(\bar{p}r^2\sin\theta\xi_\alpha\xi_\gamma\right)\right] \\ & -\frac{\partial}{\partial x_\alpha}\frac{\partial}{\partial x_\gamma}\left(\bar{p}r^2\sin\theta\xi_\alpha\frac{\partial\xi_\gamma}{\partial\phi}\right),\end{aligned} \tag{A22}$$

we find

$$\left\langle \frac{\partial}{\partial x_\alpha}\frac{\partial}{\partial x_\gamma}\left(\bar{p}r^2\sin\theta\frac{\partial\xi_\alpha}{\partial\phi}\xi_\gamma\right)\right\rangle = \left\langle \frac{\partial}{\partial x_\alpha}\frac{\partial}{\partial x_\gamma}\left(\bar{p}r^2\sin\theta\xi_\alpha\frac{\partial\xi_\gamma}{\partial\phi}\right)\right\rangle = 0, \tag{A23}$$

and hence we can omit the second term on the right-hand-side of equation (A21) to obtain

$$\left\langle \hat{r}^2 \sin\hat{\theta}\,\hat{p}\frac{\partial \xi_\gamma}{\partial x_3} K_{\gamma\alpha} \right\rangle = \left\langle r^2 \sin\theta \frac{\partial \xi_\alpha}{\partial x_3} p' \right\rangle, \tag{A24}$$

and hence

$$\langle R_{3\alpha}\rangle = \delta_{3\alpha}\left\langle J\hat{r}^2\sin\hat{\theta}\,\hat{p} - r^2\sin\theta\,\tilde{p}\right\rangle - \left\langle r^2\sin\theta\frac{\partial\xi_\alpha}{\partial x_3}p'\right\rangle. \tag{A25}$$

The meanflow equation (A18) is then reduced to

$$\bar{\rho}\frac{d\left\langle \hat{\ell}\right\rangle}{dt} = -\nabla\cdot\left\langle \boldsymbol{\xi}\frac{\partial p'}{\partial\phi}\right\rangle - \left\langle \bar{\rho}\frac{\partial\hat{\Phi}}{\partial\hat{\phi}}\right\rangle, \tag{A26}$$

where $\boldsymbol{\xi} = \xi_r\boldsymbol{e}_r + \xi_\theta\boldsymbol{e}_\theta + \xi_\phi\boldsymbol{e}_\phi$ with $\boldsymbol{e}_r$, $\boldsymbol{e}_\theta$, and $\boldsymbol{e}_\phi$ being the orthonormal vectors in the r, θ, and ϕ directions, and $\xi_r = \xi_1$, $\xi_\theta = r\xi_2$, and $\xi_\phi = r\sin\theta\xi_3$, and $\tilde{\rho} = \bar{\rho}$. Since

$$\left\langle \bar{\rho}\frac{\partial\hat{\Phi}}{\partial\hat{\phi}}\right\rangle = \left\langle \bar{\rho}\frac{\partial x_\beta}{\partial\hat{\phi}}\frac{\partial}{\partial x_\beta}\left(\bar{\Phi}+\Phi'+\boldsymbol{\xi}\cdot\nabla\bar{\Phi}\right)\right\rangle = \nabla\cdot\left\langle\bar{\rho}\boldsymbol{\xi}\frac{\partial\Phi'}{\partial\phi}\right\rangle + \left\langle\rho'\frac{\partial\Phi'}{\partial\phi}\right\rangle, \tag{A27}$$

which is correct to second order of perturbations, we obtain

$$\bar{\rho}\frac{d\left\langle \hat{\ell}\right\rangle}{dt} = -\nabla\cdot\left\langle \boldsymbol{\xi}\frac{\partial p'}{\partial\phi} + \bar{\rho}\boldsymbol{\xi}\frac{\partial\Phi'}{\partial\phi} + \frac{\nabla\Phi'}{4\pi G}\frac{\partial\Phi'}{\partial\phi}\right\rangle, \tag{A28}$$

where we have used $\partial\bar{\Phi}/\partial\phi = 0$, $\nabla^2\Phi' = 4\pi G\rho'$ and $\langle\nabla\Phi'\cdot\partial\nabla\Phi'/\partial\phi\rangle = 0$. Since the azimuthal and temporal dependence of the perturbations are assumed to be given by the factor $\exp(\mathrm{i}m\phi + \mathrm{i}\omega t)$ with m and ω being the azimuthal wavenumber and oscillation frequency, using, for example,

$$\langle\xi_\phi\xi_r\rangle = \frac{1}{2}\mathrm{Re}\left(\xi_\phi^*\xi_r\right) = \frac{1}{2}\mathrm{Re}\left(\xi_\phi\xi_r^*\right), \tag{A29}$$

we can rewrite equation (A28) as

$$\bar{\rho}\frac{d\left\langle\hat{\ell}\right\rangle}{dt} = -\frac{1}{2}\mathrm{Re}\left[\nabla\cdot\left(\boldsymbol{\xi}^*\frac{\partial p'}{\partial\phi} + \bar{\rho}\boldsymbol{\xi}^*\frac{\partial\Phi'}{\partial\phi} + \frac{\nabla\Phi'^*}{4\pi G}\frac{\partial\Phi'}{\partial\phi}\right)\right]$$

$$= \frac{m}{2}\mathrm{Im}\left[\nabla\cdot\left(\boldsymbol{\xi}^* p' + \bar{\rho}\boldsymbol{\xi}^*\Phi' + \frac{\nabla\Phi'^*}{4\pi G}\Phi'\right)\right], \tag{A30}$$

where the asterisk indicates the complex conjugation.

Integrating over a spherical surface, we obtain

$$\bar{\rho}\overline{\frac{d\left\langle\hat{\ell}\right\rangle}{dt}} = \frac{m}{2r^2}\frac{d}{dr}r^2\mathrm{Im}\left(\overline{\xi_r^* p' + \bar{\rho}\xi_r^*\Phi' + \frac{\partial\Phi'^*}{\partial r}\frac{\Phi'}{4\pi G}}\right). \tag{A31}$$

This expression (A31) is essentially the same as that used by Papaloizou & Savonije (1997) who discussed the forcing by low frequency modes tidally excited in a massive star by the orbital motion of the companion in a binary system (see also Ryu & Goodman 1992, Lin, Papaloizou, & Kley 1993). In the Cowling approximation, we obtain

$$\bar{\rho}\overline{\frac{d\left\langle\hat{\ell}\right\rangle}{dt}} = \frac{m}{2\pi r^2}\frac{dW(r)}{dr}, \tag{A32}$$

where

$$W(r) = -\pi r^2\mathrm{Im}\left(\overline{p'^*\xi_r}\right) \tag{A33}$$

may be regarded as a work function (e.g., Unno et al. 1989). We note that $dW/dr > 0$ and $dW/dr < 0$ respectively indicate the excitation and damping regions for the oscillation modes. For uniformly rotating stars, non-axisymmetric ($m \neq 0$) oscillations of rotating stars are separated into prograde and retrograde modes, and in our convention, positive (negative) m is used for retrograde (prograde) modes. From equation (A34), we find that there occurs acceleration (deceleration) of the zonal flow in the damping (excitation) regions of prograde modes ($m < 0$), while deceleration (acceleration) occurs in the damping (excitation) regions of retrograde modes ($m > 0$).

It is possible to rewrite the mean flow equation (A26) using the Reynolds stress. Substituting into equation (A26) the ϕ component of the linearized momentum equation given by

$$\frac{\partial p'}{\partial\phi} = -\bar{\rho}r\sin\theta\left[\left(\frac{\partial}{\partial t} + \frac{\bar{v}_\phi}{r\sin\theta}\frac{\partial}{\partial\phi}\right)v'_\phi + \frac{1}{r\sin\theta}\frac{\partial\bar{\ell}}{\partial r}v'_r + \frac{1}{r\sin\theta}\frac{1}{r}\frac{\partial\bar{\ell}}{\partial\theta}v'_\theta + \frac{1}{r\sin\theta}\frac{\partial\Phi'}{\partial\phi}\right], \tag{A34}$$

where $\bar{v}_\phi = r\sin\theta\Omega_1(r,\theta)$, $\bar{\ell} = \Omega r^2\sin^2\theta$, $\Omega = \Omega_1 + \Omega_c$, and using

$$\boldsymbol{v}' = \frac{\partial\boldsymbol{\xi}}{\partial t} + \Omega_1\sum_j\frac{\partial\xi_j}{\partial\phi}\boldsymbol{e}_j - r\sin\theta\left(\boldsymbol{\xi}\cdot\nabla\Omega_1\right)\boldsymbol{e}_\phi, \tag{A35}$$

we obtain after some manipulations

$$\begin{aligned}\bar{\rho}\frac{d\left\langle\hat{\ell}\right\rangle}{dt} &= -\frac{1}{r^2}\frac{\partial}{\partial r}\left\langle\bar{\rho}r^3\sin\theta v'_r\left(v'_\phi + 2\Omega f_\theta\cos\theta\xi_\theta\right)\right\rangle \\ &\quad - \frac{1}{r\sin\theta}\frac{\partial}{\partial\theta}\left\langle\bar{\rho}r\sin^2\theta v'_\theta\left(v'_\phi + 2\Omega f_r\sin\theta\xi_r\right)\right\rangle - \left\langle\rho'\frac{\partial\Phi'}{\partial\phi}\right\rangle + \frac{\partial}{\partial t}\nabla\cdot\left\langle\bar{\rho}r\sin\theta Q\boldsymbol{\xi}\right\rangle \\ &\quad - \frac{\partial}{\partial t}\left(\frac{1}{r^2}\frac{\partial}{\partial r}\left\langle\bar{\rho}r^3\sin^2\theta\Omega f_r\xi_r^2\right\rangle + \frac{1}{r\sin\theta}\frac{\partial}{\partial\theta}\left\langle\bar{\rho}r\sin^2\theta\cos\theta\Omega f_\theta\xi_\theta^2\right\rangle\right),\end{aligned} \tag{A36}$$

where

$$f_r = 1 + \frac{1}{2}\frac{\partial\ln\Omega}{\partial\ln r}, \quad f_\theta = 1 + \frac{1}{2}\frac{\partial\ln\Omega}{\partial\ln\sin\theta}, \tag{A37}$$

$$Q = v'_\phi + 2\Omega f_\theta\cos\theta\xi_\theta + 2\Omega f_r\sin\theta\xi_r. \tag{A38}$$

If we neglect the terms like $\partial\left\langle Q\boldsymbol{\xi}\right\rangle/\partial t$ and $\partial\left\langle\xi_r^2\right\rangle/\partial t$ assuming the amplitudes of the perturbations are saturated, for example, by nonlinear effects, the expression for the forcing terms in the mean flow equation (A36) with $f_\theta = 1$ reduce to that proposed by Pantillon et al. (2007) and Mathis (2009).

It is instructive to give an example of the forcing term in the meanflow equation. To see the behavior of the forcing term, we introduce the local timescale τ^{AM} defined by

$$\frac{1}{\tau^{\mathrm{AM}}} \equiv \frac{\overline{d\left\langle\ell\right\rangle/dt}}{\overline{\left\langle\ell\right\rangle}}. \tag{A39}$$



Fig. 6. $1/\tau^{\mathrm{AM}}$ for unstable $l=|m|=1$ g_{22} modes for $\Omega_c/(GM/R^3)^{1/2}=0.1$ (left panel) and an unstable r_{30} mode for $\Omega_c/(GM/R^3)^{1/2}=0.4$ (right panel) of a $4M_\odot$ main sequence model with $X=0.7$, $Z=0.02$, and $X_c=0.3646$, where the solid and dotted lines indicate retrograde and prograde modes, respectively. Note that uniform rotation is assumed. The amplitude normalization is given by $S_{l_1}(R)=1$ at the surface, and τ^{AM} is given in units of seconds.

An example of $1/\tau^{\mathrm{AM}}$ is given for low frequency modes of a $4M_\odot$ main sequence star with $X=0.7$ and $Z=0.02$ in Figure 6, where the model has been calculated with a standard stellar evolution code with the OPAL opacity (Iglesias & Rogers 1996), and we have used the method of calculation given by Lee & Saio (1993) for non-adiabatic oscillation modes of a uniformly rotating star. As the figure shows, there occurs a strong acceleration by retrograde g- and r-modes in the layers at $r/R\sim 0.95$, although the prograde g-mode contributes to deceleration of the rotation.

References

Ando H., 1983, PASJ, 35, 343
Ando H., 1986, A&A, 163, 97
Andrews D.G., McIntyre M.F., 1978a, J. Atoms. Sci., 35, 175
Andrews D.G., McIntyre M.F., 1978b, J. Fluid Mech., 89, 609
Aprilia, Lee U., Saio H., 2011, MNRAS, 412, 2265
Chandrasekhar S., 1939, An Introduction to the Study of Stellar Structure (Chicago: University of Chicago Press)
Dunkerton T., 1980, Rev. Geophys. Sp. Phys., 18, 387
Dziembowski W.A., Moskalik P., Pamyatnykh A.A., 1993, MNRAS. 265, 588
Gautschy A., Saio H., 1993, MNRAS, 267, 1071
Goldreich P., Nicholson P.D., 1989, ApJ, 342, 1075
Gough D., 1997, Nature, 388, 324
Grimshaw R., 1984, Ann. Rev. Fluid. Mech., 16, 11
Iglesias C.A., Rogers F.J., 1996, ApJ, 464, 943
Kumar P., Talon S., Zahn J.P., 1999, ApJ, 520, 859
Lee U., 2012, MNRAS, 420, 2387
Lee U., Saio H., 1993, MNRAS, 261, 415
Lee U., Saio H., Osaki Y., 1991, MNRAS, 250, 432
Lin D.N., Papaloizou J.C.B., Kley W., 1993, ApJ, 416, 689
Mathis S., 2009, A&A, 506, 811
Mathis S., Talon S., Pantillon F.P., Zahn J.P., 2008, Solar Phys., 251, 101
Neiner C., Floquet, M., Samadi, R., et al, 2012, A&A, 546, A47
Okazaki A.T., 2001, PASJ, 43, 75
Paczyński B., 1991, 370, 597
Pantillon F.P., Talon S., Charbonnel C., 2007, A&A, 474, 155
Papaloizou J.C.B., Sovonije G.J., 1997, MNRAS, 291, 651
Porter J.M., Rivinius T., 2003, PASP, 115, 1153
Press W.H., 1981, ApJ, 245, 286
Ryu D., Goodman J., 1992, ApJ, 388, 438
Schtzman E., 1993, 279, 431
Talon S., Kumar P., Zahn J.P., 2002, ApJ, 574, L175
Unno W., Osaki Y., Ando Y., Saio H., Shibahashi H., 1989, Nonradial oscillations of Stars, 2nd edn. (University of Tokyo Press)
Waters L.B.F.M., 1986, A&A, 162, 121
Willems B., van Hoolst T., Smeyers P., 2003, A&A, 397, 973
Witte M.G., Savonije G.J., 2001, A&A, 366, 840
Zahn J.P., 1975, A&A, 41, 329
Zahn J.P., 1977, A&A, 57, 383